



**PHYSICIAN'S ORDERS**

SEE MASTER ALLERGY SHEET

**SEVERE SEPSIS / SHOCK ORDERS**

Page 1 of 2

TIME ORDER NOTED

ORDERED		SEVERE SEPSIS / SHOCK ORDERS	Page 1 of 2	TIME ORDER NOTED
Date	Time			
		1. Admit to CCU: Physician:		
		2. Diagnosis: Sepsis		
		3. Consult:		
		4. Routine MICU orders		
		5. Foley catheter to gravity		
		6. Place INT, Obtain permit for placement of central line with ScvO2 monitoring		
		7. Consult Respiratory Therapy to assess and titrate O2 to maintain O2 saturations > 92%		
		<b>TESTING: IF NOT DONE IN ER:</b>		
		8. Blood culture x 2 sites, 15 minutes apart, NOW, Type/Screen for 2 units PRBCs NOW. If ScvO2 < 70 & HCT < 30% contact MD for transfusion orders.		
		9. DIC profile NOW, CMP, CBC, Magnesium, Phosphorus, Serum Lactate, ABG		
		10. Urinalysis, Urine Culture/Sensitivity, Sputum for gram stain/C & S		
		11. Portable CXR		
		<b>MEDICATIONS:</b>		
		12. STAT ANTIBIOTICS:		
		13. Accuchecks every 4 hours		
		14. If blood sugar > 125, start Regular Insulin per protocol: <input type="checkbox"/> Renal Failure (ESRD)		
		<input type="checkbox"/> Standard <input type="checkbox"/> Intensive		
		15. Acetaminophen (Tylenol) 650 mg po or per rectum every 4 hours PRN Temp > _____ (max of 4 GM/day)		
		16. Volume replacement via central line and continuous CVP measure		
		17. Give Normal Saline 500 ml bolus every 30 minutes until CVP goal met (CVP goal is greater than 8mm Hg non vent; & CVP greater than 12mm Hg on vent)		
		18. If MAP remains < 65, start:		
		<input type="checkbox"/> Norepinephrine (Levophed) 2 mcg/min IV, titrate to keep MAP > 65, max dose 20		
		<input type="checkbox"/> If no response to Norepinephrine, start Dopamine 5 mcg/kg/min IV, titrate to keep MAP > 65, max dose 20 mcg/kg/min		
		<input type="checkbox"/> If MAP < 65 and/or ScvO2 < 70 on high dose of Dopamine & Norepinephrine – Add Vasopressin – Start 0.02 units/minute IV		
		19. If ScvO2 < 70 and HCT > 30% contact MD for inotropic agent		
		20. <input type="checkbox"/> Cosyntropin Stimulation Test:		
		Serum Cortisol level now, Give cosyntropin 250 mcg IV immediately after level drawn		
		Repeat serum cortisol level 30 minutes after cosyntropin given		
		Physician Signature: _____		
		Date/Time: _____		

▼ Patient Label ▼



**SEVERE SEPSIS / SHOCK ORDERS**